







































(Pseudo Kaposi)

Consider

Consider the following: **nasopharyngeal carcinoma** (in children)

Always Rule Out: **Alveolar rhabdomyosarcoma** (in children)
Angiosarcoma (in the liver, soft tissue malignancies and leukemias)

Away from Rule: The following are considered normal malignancies and leukemias

(continued)

Discriminate between the two methods of involvement

Disseminated *M. neoformans* involvement in active-antiretroviral therapy-naïve HIV-infected patients

alone system. The results of the study are as follows: The active carbon adsorption capacity of the

[illegible]

ANTISEPTIC: Antiseptics are used to disinfect the skin with a preselection for head/neck

Amplification of *hsp90* mRNA in cultured human synovial cells grown in the presence of interleukin-1 α under basal and high concentrations of dexamethasone. The cells were grown in the presence of 1 ng/ml dexamethasone or 100 ng/ml dexamethasone for 48 h before the addition of 1 ng/ml interleukin-1 α for 24 h. The cells were then harvested and total RNA was extracted. The amount of *hsp90* mRNA was determined by Northern blot analysis. The results are expressed as the mean \pm SD of three independent experiments. **P* < 0.05 compared to the control.

evaluation of the impact of the proposed changes on the environment, the proposed changes to the existing project are not expected to have any significant adverse effects on the environment.